



04.1142.5017

MFM GRAZER R (Rumensin) MINERAL

MARTINDALE GRAZER (R)

MEDICATED

For Beef Cattle on Pasture

For increased rate of weight gain in pasture cattle (slaughter, stocker, and feeder cattle, and dairy and beef replacement heifers).

DO NOT FEED UNDILUTED

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 1620 g/ton

GUARANTEED ANALYSIS

| | |
|--|---------|
| Calcium (Ca), minimum | 10.30% |
| Calcium (Ca), maximum | 12.30% |
| Phosphorus (P), minimum | 3.50% |
| Salt (NaCl), minimum | 14.60% |
| Salt (NaCl), maximum | 17.50% |
| Potassium (K), minimum | 0.60% |
| Copper (Cu), minimum ppm | 830 |
| Selenium (Se), minimum ppm | 26.40 |
| Zinc (Zn), minimum ppm | 2,000 |
| Vitamin A, minimum IU per lb | 300,000 |
| Vitamin D ₃ , minimum IU per lb | 30,000 |
| Vitamin E, minimum IU per lb | 100 |

INGREDIENTS

Monocalcium Phosphate, Dicalcium Phosphate, Calcium Carbonate, Salt, Molasses Products, Corn Distillers Dried Grains with Solubles, Cottonseed Meal, Vitamin A Acetate, Vitamin D₃ Supplement, Vitamin E Supplement, Cobalt Carbonate, Copper Sulfate, Calcium Iodate, Ferrous Carbonate, Manganous Oxide, Sodium Selenite, Zinc Oxide, Blended Vegetable Fats (preserved with BHA and BHT), Artificial Flavors

FEEDING DIRECTIONS

Thoroughly mix Vigortone Southwest Grazer (R) Medicated with grain or roughage before feeding.

Feed 1 to 3.95 ounces of Vigortone Southwest Grazer (R) Medicated mixed in 1 lb or more of feed per head daily to supply 50 to 200 mg of monensin per head per day. During the first 5 days of feeding, feed 1 to 1.97 ounces of Vigortone Southwest Grazer (R) Medicated so cattle receive no more than 100 mg of monensin per head per day.

CAUTION: Do not allow horses or other equines access to formulations containing monensin. Ingestion of monensin by equines has been fatal.

Monensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions.

Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats.

Must be thoroughly mixed in feeds before use.

Do not feed undiluted.

Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result.

Do not feed to lactating dairy cows. Do not feed to lactating goats.

A withdrawal period has not been established for monensin in peruminating calves. Do not use in calves to be processed for veal.

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

Prt'd. 10/02

Net Weight 50 lb (22.68 kg)

(800)366-9584
www.martindalefeed.com