



04.3333.9620



20% All Natural Compressed Block

For Beef Cattle and Horses on Pasture

Guaranteed Analysis:

Crude Protein (Min)	20.00%
Crude Fat (Min)	5.00%
Crude Fiber (Max)	8.00%
Calcium (Min)	4.00%
Calcium (Max)	5.00%
Phosphorus (Min)	1.00%
Salt (Min)	2.00%
Salt (Max)	2.50%
Magnesium (Min)	3.00%
Copper (Min)	100 ppm
Manganese (Min)	400 ppm
Selenium (Min)	3.3 ppm
Zinc (Min)	300 ppm
Vitamin A (Min)	25,000 IU/lb
Vitamin D-3 (Min)	2,500 IU/lb
Vitamin E (Min)	25 IU/lb

CAUTION: USE AS DIRECTED. Consumption of selenium should not exceed 3 mg per head per day.

WARNING: This product, which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.

NET WEIGHT 200 LB. (90.7 KG)

Ingredients:

Processed Grain By-Products, Molasses Products, Condensed Fermented Corn Extractives, Magnesium Oxide, Calcium Hydroxide, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Plant Protein Products, Calcium Carbonate, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Chloride, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydrochloride, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement and Vitamin E Supplement.

INTRODUCTORY PERIOD

Following a 7 to 10 day acclimation period, average consumption should be 1 to 2 pounds per head per day for mature cattle and horses with unrestricted forage supplies. Position the blocks 50 to 75 feet from the loafing, feeding and watering areas during the first week of feeding. Some repositioning of blocks may be necessary as animals become acclimated to the presence of the blocks. Move blocks closer to increase consumption; further away to decrease consumption. If overconsumption occurs for more than two weeks and repositioning of blocks does not correct the situation, remove blocks and evaluate the overall feeding program. **DO NOT ALLOW STARVED ANIMALS FREE ACCESS TO THIS BLOCK.**

FEEDING RECOMMENDATIONS

1. Provide 1 block per 10 to 20 head. All animals should have equal access to blocks. Timid and smaller animals need their share.
2. Place blocks where animals congregate. Locate blocks throughout the pasture at loafing, grazing, feeding and watering areas. Adequate distance must be maintained between blocks to minimize crowding - 10 feet minimum.
3. Feed blocks continuously along with a plentiful source of forage and clean, fresh water. Animals should not run out of blocks. When a block is two-thirds consumed, provide a new block near it.
4. Cattle and horses should consume an average of 1 to 2 pounds per head daily based on a 1,000-pound animal.
5. Consumption of this block may vary depending upon animal body condition, quality and quantity of forages, seasonal weather conditions, and most importantly, feeding locations of blocks with respect to loafing, grazing, feeding and watering areas.
6. If additional mineral and vitamin fortification is desired, provide an appropriate SWEETLIX® free choice mineral supplement in addition to this supplement.

For additional information, contact your SWEETLIX® representative.

9/10

Manufactured By:
Ridley Block Operations
P.O. Box 8500, Mankato, MN 56002
www.sweetlix.com



27976

(800)366-9584
www.martindalefeed.com