



04.3316.4020

SWEETLIX HORSE/CATTLE BLOCK



Rabon[®] Molasses Pressed Block

A Free Choice Supplement Block for Cattle and Horses

For control of fecal flies in manure of treated cattle and horses. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle. Prevents development of house flies and stable flies in the manure of treated horses.

ACTIVE INGREDIENT:

Tetrachloro-phos: (Z)-2-Chloro-1-(2,4,5-Trichlorophenyl) vinyl dimethyl phosphate*	0.494%
Other Ingredients**	99.506%
	100.000%

*Each lb. contains 2.24 g. Rabon[®]
 **Refers only to ingredients which are not inert.

GUARANTEED ANALYSIS

Calcium (Min)	4.50%
Calcium (Max)	5.50%
Phosphorus (Min)	4.00%
Salt (Min)	15.00%
Salt (Max)	18.00%
Magnesium (Min)	1.00%
Potassium (Min)	2.00%
Iodine (Min)	90 ppm
Copper (Min)	300 ppm
Manganese (Min)	800 ppm
Selenium (Min)	21 ppm
Zinc (Min)	1,200 ppm
Vitamin A (Min)	100,000 IU/lb.
Vitamin D-3 (Min)	25,000 IU/lb.
Vitamin E (Min)	100 IU/lb.

FEED INGREDIENTS

Cane Molasses, Soybean Hulls, Dicalcium Phosphate, Monocalcium Phosphate, Salt, Potassium Chloride, Magnesium Limestone, Calcium Carbonate, Magnesium Oxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Calcium Iodide, Cobalt Sulfate, Zinc Sulfate, Ferrous Sulfate, Zinc Oxide, Manganese Oxide, Manganese Sulfate, Copper Sulfate, Sodium Selenite, Yeast Culture, Lignin Sulfonate and Petrolatum.

Keep Out of Reach of Children

CAUTION

EPA Reg. No. 73600-1 • EPA Est. # 73600-TX-01

Manufactured By Sweetlix, PO Box 8500, Mankato, MN 56002
 Contact us at: 1-875-SWEETLIX • www.sweetlix.com

First Aid	
IF SWALLOWED	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.
IF ON SKIN	Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF IN EYES	Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or physician for treatment advice.
HOT LINE NUMBER: In case of emergency, contact Sweetlix (800) 325-1486.	
NOTE TO PHYSICIANS AND VETERINARIANS: Symptomatology: Signs of organophosphate toxicity are those of cholinergic over stimulation; hypersalivation, excessive urination, diarrhea and dyspnea (respiratory distress, collapse and death from respiratory muscle paralysis). Treatment: Withdraw product immediately. Clinical includes 3 categories: muscarinic blocking agents (atropine sulfate), (2) cholinesterase reactivators (atropine and 2-PAM, pralidoxime chloride) and (3) emetics (i.e. syrup of ipecac), cathartics (i.e. vegetable oil) and absorbents (i.e. charcoal slurry).	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

USER SAFETY RECOMMENDATIONS

PERSONAL PROTECTIVE EQUIPMENT (PPE): Long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS: This compound is toxic to fish.

DIRECTIONS FOR USE

If is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: Not to be used on horses destined for slaughter.

PRECAUTIONS FOR HORSE USE: This product contains tetrachloro-phos, which is a cholinesterase inhibitor. The most frequently reported clinical signs of cholinesterase inhibition in the horse are abdominal

pain, lethargy, sweating, tearing and excessive salivation. If these signs are seen in horses, consult your veterinarian immediately. Do not use this product simultaneously or within a week before or after treatment with cholinesterase inhibiting drugs, pesticides or chemicals. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals. This product has not been tested in breeding horses.

GENERAL INSTRUCTIONS

1. Feed 1 block per 5 head of cattle or horses. This allows all animals equal access to blocks. Feed blocks at the rate of 0.5 oz. per 100 lb. of bodyweight per day. This intake will supply 0.07g. of larvicide per 100 lb. of bodyweight per day (2.1 grams per 100 lbs. bodyweight per month).
2. Place blocks where animals congregate. Locate blocks throughout pasture, placing them near water and loafing areas. Maintain adequate distance between blocks (10 ft. minimum).
3. Do not allow animals to run out of blocks. When a block is half-consumed, place a new block near it. Place very small pieces in troughs or feed pans.
4. Animals will consume about 5 oz. per head daily. When animals consume the block at the recommended rate, it will supply 3 mg of selenium.
5. Blocks can be fed to animals in confinement. Exact location of blocks to obtain desired consumption will vary between confinement feeding programs.
6. Feed as the only FREE CHOICE source of salt, other minerals and vitamins.
7. SWEETLIX[®] Rabon[®] Molasses Pressed Block prevents the development of fecal flies in the manure of treated cattle and horses, but is not effective against existing adult flies. It may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which can breed not only in animal manure but in other decaying matter on the premises. In order to achieve optimum fly control, SWEETLIX[®] Rabon[®] Molasses Pressed Block should be used in conjunction with other good management and sanitation practices. Start feeding SWEETLIX[®] Rabon[®] Molasses Pressed Block early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. SWEETLIX[®] Rabon[®] Molasses Pressed Block may be fed to cattle up to slaughter.

8. Follow feeding directions. Consumption of selenium should not exceed 3 mg per head daily.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.

Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and Local Authorities, by burning. If burned, stay out of smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

For additional information, please contact your SWEETLIX representative. Rabon[®] is a registered trademark of KMG Chemicals, Houston, TX

NET WEIGHT 40 LBS. (18.18 KG)